

Interference and Distinguishability in Quantum Mechanics

Jos Uffink and Jan Hilgevoord

April, 1988

Abstract

Quantitative measures are introduced for the indistinguishability U of two quantum states in a given measurement and the amount of interference I observable in this measurement. It is shown that these measures obey an inequality $U \geq I$ which can be seen as an exact formulation of Bohr's claim that one cannot distinguish between two possible paths of a particle while maintaining an interference phenomenon. This formulation is applied to a neutron interferometer experiment of Badurek e.a. It is shown that the formulation is stronger than an argument based on an uncertainty relation for phase and photon number considered by these authors.

A recent experiment in neutron interferometry [1] can be seen as a realisation of the double-slit thought experiment discussed by Einstein and Bohr. In this famous discussion, Bohr argued that one cannot distinguish between two possible paths of a particle while preserving an interference phenomenon. To reach this conclusion Bohr applied the uncertainty relation for position and momentum in a somewhat informal way. However, it has been shown that the Heisenberg uncertainty relation by itself is not strong enough to justify Bohr's claim [2].

The discussion has been revived in the light of the new neutron experiments. In ref. 1 an argument is presented supporting Bohr's claim, based on an uncertainty relation for the phase and photon number of the electromagnetic field. The validity of this explanation was subsequently disputed [3], because of the dubious theoretical status of the phase-number uncertainty relations.

This raises the question whether it is possible to give a direct quantitative formulation of Bohr's claim, without recourse to the uncertainty relations. Work in this direction has been done by Wootters and Zurek[4]. Here we propose an alternative formulation that seems particularly apt for the interferometer experiments.

The main obstacle for a direct formulation of Bohr's claim is the problem of choosing a quantitative measure for the extent to which the two paths are distinguishable in a given experiment. Let ψ_1 and ψ_2 denote orthogonal quantum states that represent possible paths of a particle. A measurement performed on this particle may be described by a complete set of orthogonal projection operators $\{D_k\}$, where k denotes a possible outcome of the measurement. The hypotheses that the particle traveled either one of two paths then provide two probability distributions, viz.

$$\begin{aligned} p_k &= \langle \psi_1 | D_k | \psi_1 \rangle, \\ q_k &= \langle \psi_2 | D_k | \psi_2 \rangle. \end{aligned} \tag{1}$$

The two paths may be said to be discriminated if, from the observed outcome of the measurement one can decide between these two hypotheses. Thus, the problem of distinguishing between two paths can be seen as a special case of the general classical problem of discriminating between two statistical hypotheses.

A solution to this problem depends, of course, on the distributions p and q , but also on the observed outcome. However, independently of the latter, one can indicate whether a discriminative answer is likely. There are two extreme cases. (i): $p_k q_k = 0$ for all k , i.e. every outcome that has positive probability according to one hypothesis is impossible according to the other. In this case a single observation will suffice for complete discrimination. (ii): $p_k = q_k$ for all k . In this case, no number of observations can discriminate between p and q . In all other cases an incomplete discrimination is to be expected.

One is tempted to define a ‘degree of indistinguishability’, reflecting the expected lack of discrimination between p and q . For this purpose we choose

$$U(p, q) = \sum_k \sqrt{p_k q_k}. \quad (2)$$

The significance of this expression in statistical theory has been studied by Bhattacharyya [5], Rao [6] and Wootters [7]. Loosely speaking, a value of $U(p, q)$ close to unity indicates that even when one of these distributions is ‘true’ a typical outcome of the experiment will not allow one to infer accurately which of them is true.

As an illustration we compare the expression proposed above with a more familiar statistical approach to the discrimination problem, the theory of hypothesis testing [8]. A ‘test’ for two statistical hypotheses is a procedure by which one hypothesis is either accepted or rejected in favor of the alternative hypothesis. In this approach the quality of the test is judged by the so-called errors of the first and second kind, $1 - \alpha$ and $1 - \beta$; i.e., respectively, the probability of rejecting the first hypothesis when it is true, and the probability of accepting it when its alternative is true. A Neyman-Pearson (NP) test’ is designed to minimize these two errors, and always obeys $(1 - \alpha) + (1 - \beta) \leq 1$. One can show that for all NP tests of the distributions p and q :

$$\begin{aligned} (1 - \alpha) + (1 - \beta) &\geq 1 - \sqrt{1 - U^2}, \\ (1 - \alpha)(1 - \beta) &\leq \frac{1}{4} U^2. \end{aligned} \quad (3)$$

Thus, a large degree of indistinguishability implies that the sum of the two kinds of error in a NP test is close to its upper bound 1, whereas a small value of $U(p, q)$ implies that any NP test of p and q is ‘good’ in the sense that the product of the errors is small.

In agreement with (2) we define the degree of indistinguishability of the quantum states ψ_1 and ψ_2 for the measurement $\{D_k\}$ as

$$U_{\{D_k\}}(\psi_1, \psi_2) = \sum_k \sqrt{\langle \psi_1 | D_k | \psi_1 \rangle \langle \psi_2 | D_k | \psi_2 \rangle}. \quad (4)$$

Thus even when it is assumed that the particle did travel one of the paths, the outcome of the measurement will only enable us to determine this path if U is small.¹ One may easily show that for given ψ_1 and ψ_2 the expressions (4) is non-decreasing when the projections are resolved into lower-dimensional projections, as

¹(Note added:) The meaning of the indistinguishability of two hypotheses is explained here in terms of the assumption that the particle does actually travel a definite path. One may object that, according to Bohr, this is not the case in a interference experiment. Indeed, in that case, the true state is not ψ_1 nor ψ_2 but a superposition of both. However, the assumption is made here only for the purpose of elucidating the meaning of ‘indistinguishability’. The point is that depending on whether $U_{\{D_k\}}(\psi_1, \psi_2)$ is large (or small), the two states are hard (or easy) to tell apart. Once this is accepted, the question whether $U_{\{D_k\}}(\psi_1, \psi_2)$ is indeed small or large is of course completely independent of the question whether ψ_1 , ψ_2 or any other state is actually prepared in the experiment.

is intuitively reasonable. Maximal distinguishability is reached when the set $\{D_k\}$ includes $|\psi_1\rangle\langle\psi_1|$ or $|\psi_2\rangle\langle\psi_2|$.

We now turn to the notion of interference. The state of a particle emerging from an interferometer may be regarded as a normalized superposition of ψ_1 and ψ_2 , $\psi = c_1\psi_1 + c_2\psi_2$, with $|c_1|^2 + |c_2|^2 = 1$. The probability that the measurement $\{D_k\}$ yields an outcome k is then

$$P_k = |c_1|^2 p_k + |c_2|^2 q_k + i_k, \quad (5)$$

where p_k and q_k are given by (1), and

$$i_k = 2 \operatorname{Re} c_1^* c_2 \langle \psi_1 | D_k | \psi_2 \rangle \quad (6)$$

is the so-called interference term. The maximum value of i_k for all choices of c_1 and c_2 is

$$I_k = |\langle \psi_1 | D_k | \psi_2 \rangle|.$$

We define the interference power of ψ_1 and ψ_2 for this measurement as

$$I_{\{D_k\}}(\psi_1, \psi_2) = \sum_k I_k = \sum_k |\langle \psi_1 | D_k | \psi_2 \rangle|. \quad (7)$$

Let us see how this relates to other familiar notions of interference strength. In many experiments a variable phase shift χ between the two paths is introduced. In that case it is convenient to put $\psi_2(\chi) = e^{i\chi}\psi_2$. Let us further take $c_1 = c_2 = 1/\sqrt{2}$. P_k then oscillates as a function of χ :

$$P_k(\chi) = \frac{1}{2}(p_k + q_k) + |\langle \psi_1 | D_k | \psi_2 \rangle| \cos \chi.$$

A well-known measure for the amount of interference in this situation is the Michelson fringe visibility

$$V_k = \frac{P_{\max} - P_{\min}}{P_{\max} + P_{\min}},$$

where $P_{\max}$ and $P_{\min}$ denote adjacent maximal and minimal values of $P_k(\chi)$. This gives

$$V_k = 2|\langle \psi_1 | D_k | \psi_2 \rangle|/(p_k + q_k),$$

or

$$I_{\{D_k\}} = \sum_k V_k \frac{1}{2}(p_k + q_k).$$

Thus, the interference power is just the average visibility over all possible outcomes, weighted by the mean probability $\frac{1}{2}(p_k + q_k)$.

For given ψ_1 and ψ_2 , $I_{\{D_k\}}$ is non-decreasing when the projections are resolved into lower-dimensional projections, so that a more resolving measurement will in general show more interference.

Now, let us compare the interference power (7) with the degree of indistinguishability (4). By the Schwartz inequality, one finds

$$U_{\{D_k\}}(\psi_1, \psi_2) \geq I_{\{D_k\}}(\psi_1, \psi_2). \quad (8)$$

Thus the appearance of a pronounced interference phenomenon in an experiment is incompatible with the requirement that the interfering states are distinguishable in that experiment. This can be regarded as a quantitative expression of Bohr's claim. Equality in (8) occurs when all projections D_k are one-dimensional.

In relation (8) the degree of indistinguishability and interference power are compared for one and the same experiment. What if the interference is observed in

one measurement, $\{D_k\}$, and one attempts to distinguish the states by means of another measurement, described by the set $\{D'_l\}$? Actually this will not improve the situation, as long as the two measurements are compatible. In that case, the product of two projections, $D_{kl} = D_k D'_l$, is also an orthogonal projection, and D_{kl} is a resolution of $\{D_k\}$ as well as $\{D'_l\}$, so that

$$U_{\{D'_l\}} \geq U_{\{D_{kl}\}} \geq I_{\{D_{kl}\}} \geq I_{\{D_k\}}. \quad (9)$$

We now apply the ideas discussed above to the neutron interferometer. In the interferometer an incident neutron beam is coherently split in two partial beams, 1 and 2. A phase shift may be produced by placing a piece of material in one of them. Next the two beams are again coherently split and pairwise superposed, so that the neutrons emerge in two final beams, A and B . The emerging neutron state may then be written as a superposition of orthogonal parts corresponding to the two paths.

$$\phi^0 = \frac{1}{\sqrt{2}}(\phi_1^0 + e^{i\chi}\phi_2^0), \quad (10)$$

where χ is the variable phase shift. In the ideal case, each of the paths corresponds with equal amplitude to the emerging beams,

$$\begin{aligned} \phi_1^0 &= \phi_{1A}^0 + \phi_{1B}^0, \\ \phi_2^0 &= \phi_{2A}^0 + \phi_{2B}^0, \end{aligned} \quad (11)$$

and

$$\phi_{1A}^0 = \phi_{2A}^0, \quad \phi_{1B}^0 = -\phi_{2B}^0. \quad (12)$$

Thus, for $\chi = 0$, all neutrons are found in beam A , whereas for a phase shift $\chi = \pi$ all neutrons are found in beam B . This conforms to maximal interference as a function of the phase shift.

In the experiment of ref. 1 the incident beam is prepared with spin polarized along the z -axis. One of the paths is lead through a magnetic coil which reverses the spin of the passing neutrons. In the emerging beams spin analyzers and detectors are placed, and the time-dependent intensity is recorded. When the analyzers are turned to to analyze spin in the y -direction, a periodical interference pattern in the intensity is observed as a function of the phase shift.

The question is now whether one can infer, without disturbing the interference, which path each neutron traveled through the interferometer by means of an extra measurement on the spin flipping coil. The idea behind this is that the interaction of the neutron with the magnetic field of the coil involves the exchange of a photon. Hence if a measurement of the photon number of this field is made, one might hope to detect whether or not such an interaction has taken place.

Let the spin flipper be placed in path 2. The combined final state of neutron and magnetic field may be taken as

$$\psi = \frac{1}{\sqrt{2}}(\psi_1 + e^{i\chi}\psi_2), \quad (13)$$

with

$$\psi_1 = \phi_1^0 \xi, \quad (14)$$

where ξ is the final state of the magnetic field when no interaction occurs, and ψ_2 is the combined final state of neutron and magnetic field when the interaction does take place. Introducing the neutron eigenstates for the z -component of the spin, $|\pm\rangle$, and photon number eigenstates $|n\rangle$, we may write:

$$\phi_1^0 = |+\rangle\langle+|\phi_1^0\rangle, \quad (15)$$

$$\xi = \sum_n |n\rangle\langle n|\xi\rangle. \quad (16)$$

The spin flipper is assumed to be efficient, i.e. it fully reverses the spin of all passing neutrons. Further, we assume that the interaction does not alter the spatial wave function of the neutron. ψ_2 may then be written as

$$\psi_2 = \phi_2 \xi_2, \quad (17)$$

with

$$\phi_2 = e^{i\omega t} |-\rangle \langle + | \phi_2^0, \quad (18)$$

$$\xi_2 = \sum_n e^{-i\omega t} |n+1\rangle \langle n | \xi, \quad (19)$$

where ω is the frequency of the electromagnetic field.

Suppose now that a measurement is made of the intensity in the emerging neutron beams with spin analyzed in the y -direction at all positions x along the beams at a fixed time. It is easy to see that this procedure, and the actual experiment, in which a time-dependent observation is made at a fixed position, are equivalent in the quasi-monochromatic approximation for the neutron wave packet.

The interference power of the states is (14) and (17) for this experiment is:

$$I_{\{D_{xs}\}} = |\langle \xi | \xi_2 \rangle| \sum_s \int dx |\langle \phi_1^0 | D_{xs} | \phi_2 \rangle|, \quad (20)$$

where the integral is to be performed over both emerging beams. Using the fact that the projections D_{xs} can be factorized, $D_{xs} = D_x D_s$, with

$$\begin{aligned} D_{s=+\frac{1}{2}} &= (|+\rangle + |-\rangle)(\langle +| + \langle -|), \\ D_{s=-\frac{1}{2}} &= (|+\rangle - |-\rangle)(\langle +| - \langle -|). \end{aligned}$$

We obtain from (11) and (12)

$$I_{\{D_{xs}\}} = \left| \sum_n \langle \xi | n \rangle \langle n+1 | \xi \rangle \right|. \quad (21)$$

Let us now ask whether the passage of a neutron through the spin flipper, and the associated photon exchange, can be detected by a measurement of the photon number. The two states to be distinguished are (14) and (17). For the measurement of the photon number, represented by the projections $D_n = |n\rangle \langle n|$, the degree of indistinguishability (4) is

$$U_{\{D_n\}}(\psi_1, \psi_2) = \sum_n \sqrt{|\langle \xi | n \rangle \langle n+1 | \xi \rangle|}. \quad (22)$$

Thus, the extent to which one can determine whether a photon exchange has occurred, depends on the extent to which consecutive pairs of photon numbers are represented in the distribution $|\langle n | \xi \rangle|^2$. It is obvious from (21) and (22) that

$$U_{\{D_n\}}(\psi_1, \psi_2) \geq I_{\{D_{xs}\}}(\psi_1, \psi_2), \quad (23)$$

in agreement with (9). This shows that the conditions under which a measurement on the spin flipper allows the distinction of the two paths exclude the conditions under which interference occurs.

Finally, we compare the formulation of Bohr's claim given here with an approach employing an uncertainty relation for the phase and photon number. A satisfactory

description of the phase of an electromagnetic fields in quantum theory is that in terms of the exponential phase operator $e^{i\phi}$ defined by Lévy-Leblond[9],²

$$e^{i\phi} = \sum_n |n\rangle\langle n+1|.$$

Note that the expectation value of this operator appears in (21). An appropriate definition for the phase uncertainty, as discussed in ref. 9, is

$$(\Delta\phi)^2 = 1 - |\langle\xi|e^{i\phi}|\xi\rangle| - |\langle 0|\xi\rangle|^2 \quad (24)$$

where $|\langle 0|\xi\rangle|^2$ is the probability to find zero photons in the magnetic field. Under experimental conditions this term may be neglected. Therefore, in order to have an appreciable interference power (21), it is necessary for the magnetic field that $\Delta\phi \ll 1$. We can now employ the uncertainty relation for phase and photon number given in ref. 9:

$$(\Delta n)^2((\Delta\phi)^2 - \frac{1}{2}|\langle 0|\xi\rangle|^2) \geq \frac{1}{4}(1 - \Delta\phi)^2 - |\langle 0|\xi\rangle|^2 \quad (25)$$

where $(\Delta n)^2 = \langle n^2 \rangle - \langle n \rangle^2$ is the standard deviation of the photon number in the state ξ . It then follows from condition (25) that

$$\Delta n \gg \frac{1}{2}. \quad (26)$$

However, this conclusion by itself is not sufficient to exclude the distinguishability of the two paths. For example, if the state of the photon field were such that

$$|\langle n|\xi\rangle|^2 = \frac{1}{2}(\delta_{nn_0} + \delta_{nn_1}), \quad |n_0 - n_1| \gg 1, \quad (27)$$

(where δ denotes the Kronecker delta) the photon exchange could still be detected with complete certainty from a measurement of the photon number, without violating (27). However, the fact that such a state is inconsistent with condition (25) follows directly from relation (23).

We conclude that the formulation of Bohr's claim considered above is stronger than one based on an certainty relation of the Heisenberg type.

References

- [1] G. Badurek, H. Rauch, J. Summhammer, *Phys. Rev. Lett* **51** (1983) 1015.
- [2] J. Uffink and J. Hilgevoord *Found. Phys.* **15** (1985) 925.
- [3] C. Dewdney, A. Garuccio, Ph. Gueret, A. Kyprianides and J.P.Vigier *Found. Phys.* (1985) 1035.
- [4] W.K. Wootters and W. Zurek, *Phys. Rev. D* **19** (1979) 473.
- [5] A. Bhattacharyya, *Bull. Calcutta Math. Soc.* **35** (1943) 99.
- [6] C.R. Rao, *Bull. Calcutta Math. Soc.* **37** (1945) 81.
- [7] W.K. Wootters *Phys. Rev. D* **23** (1981) 357.
- [8] M.G. Kendall and A. Stuart *The Advanced Theory of Statistics*, Vol. 2. (Griffin, London, 1946).
- [9] J-M. Lévy-Leblond, *Ann. Phys.* **101** (1976) 319.

²(Note added:) The notation ' $e^{i\phi}$ ' for this operator is merely formal and should not be mistaken as implying that it is an exponent of of some self-adjoint operator ϕ . In fact, $e^{i\phi}$ is not even unitary.